


Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 5 of 62

**COMPLETE LIST OF CLAIMS THAT ARE OR HAVE BEEN BEFORE THE
OFFICE AFTER ENTRANCE OF THE AMENDMENTS MADE HEREIN**
(See next page)

1. (CURRENTLY AMENDED) A compound of Formula (I):



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Q is $-(CR^7R^{7a})_m-R^4$,
 $-(CR^7R^{7a})_n-S-R^4$,
 $-(CR^7R^{7a})_n-O-R^4$,
 $-(CR^7R^{7a})_m-N(R^{7b})-R^4$,
 $-(CR^7R^{7a})_n-S(=O)-R^4$,
 $-(CR^7R^{7a})_n-S(=O)_2-R^4$, or
 $-(CR^7R^{7a})_n-C(=O)-R^4$;

provided when n is 0, then R⁴ is not H;

m is 1, 2, or 3;

n is 0, 1, or 2;

R⁴ is H,

C₁-C₈ alkyl substituted with 0-3 R^{4a},
 C₂-C₈ alkenyl substituted with 0-3 R^{4a},
 C₂-C₈ alkynyl substituted with 0-3 R^{4a},
 C₃-C₁₀ carbocycle substituted with 0-3 R^{4b},
 C₆-C₁₀ aryl substituted with 0-3 R^{4b}, or

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 6 of 62

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b};

R^{4a}, at each occurrence, is independently selected from is H, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, OR^{14a}, OR²², SR²², C(=O)OR²², NR²¹R²², S(=O)R²², S(=O)₂R²², C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-, C₃-C₁₀ carbocycle substituted with 0-3 R^{4b}, C₆-C₁₀ aryl substituted with 0-3 R^{4b}, and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b};

R^{4b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁵ is H;

C₁-C₆ alkyl substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b};
C₂-C₆ alkynyl substituted with 0-3 R^{5b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶;
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 7 of 62

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁶ is H;

C₁-C₆ alkyl substituted with 0-3 R^{6a};

C₃-C₁₀ carbocycle substituted with 0-3 R^{6b}; or

C₆-C₁₀ aryl substituted with 0-3 R^{6b};

R^{6a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, aryl or CF₃;


R^{6b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy;

R⁷, at each occurrence, is independently H or C₁-C₄ alkyl;

R^{7a}, at each occurrence, is independently H or C₁-C₄ alkyl;

R^{7b} is H or C₁-C₄ alkyl;

Ring B is



a 7-membered lactam;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 8 of 62

~~wherein the lactam is saturated, partially saturated or unsaturated;
wherein each additional lactam carbon is substituted with 0-2 R¹¹; and,
optionally, the lactam contains a heteroatom selected from O, S, S(=O), S(=O)₂,
N=, NH, and N(R¹⁰);~~

~~additionally, two R¹¹ substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R¹³;~~

~~additionally, two R¹¹ substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical; wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R¹³;~~

~~additionally, two R¹¹ substituents on the same or adjacent carbon atoms may be combined to form a C₃-C₆ carbocycle substituted with 0-3 R¹³;~~

~~R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹,
S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷,~~

~~C₁-C₆ alkyl optionally substituted with 0-3 R^{10a},~~

~~C₆-C₁₀ aryl substituted with 0-4 R^{10b},~~

~~C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or~~

~~5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{10b};~~

~~R^{10a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃, or aryl substituted with 0-4 R^{10b};~~

~~R^{10b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl S;~~

~~R¹¹, at each occurrence, is independently selected from~~

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 9 of 62

~~H, C₁-C₄ alkoxy, Cl, F, Br, I, -O, CN, NO₂, NR¹⁸R¹⁹, C(=O)R¹⁷, C(=O)OR¹⁷,
C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃,
C₁-C₆ alkyl optionally substituted with 0-3 R^{11a},
C₆-C₁₀ aryl substituted with 0-3 R^{11b},
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}, or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen,
oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with
0-3 R^{11b},~~

~~R^{11a}, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, -O, CN, NO₂, NR¹⁵R¹⁶, CF₃,
phenyl substituted with 0-3 R^{11b},
C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}, and
5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen,
oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-
3 R^{11b},~~

~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂,
NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl S,~~

W is a bond or -(CR⁸R^{8a})_p;

p is 0, 1, 2, 3, or 4;

R⁸ and R^{8a}, at each occurrence, are independently selected from H, F, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl and C₃-C₈ cycloalkyl;

X is a bond;

C₆-C₁₀ aryl substituted with 0-3 R^{Xb},
C₃-C₁₀ carbocycle substituted with 0-3 R^{Xb}, or
5 to 10 membered heterocycle substituted with 0-2 R^{Xb},

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 10 of 62

R^{Xb} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ halothioalkoxy;

Y is a bond or $-(CR^9R^{9a})_t-V-(CR^9R^{9a})_u-$;

t is 0, 1, or 2;

u is 0, 1, or 2;

R⁹ and R^{9a}, at each occurrence, are independently selected from H, F, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

V is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-N(R^{19})-$, $-C(=O)NR^{19b}-$, $-NR^{19b}C(=O)-$, $-NR^{19b}S(=O)_2-$, $-S(=O)_2NR^{19b}-$, $-NR^{19b}S(=O)-$, $-S(=O)NR^{19b}-$, $-C(=O)O-$, or $-OC(=O)-$;

Z is H;

C₁-C₈ alkyl substituted with 0-3 R^{12a};

C₂-C₆ alkenyl substituted with 0-3 R^{12a};

C₂-C₆ alkynyl substituted with 0-3 R^{12a};

~~C₆-C₁₀~~ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, $-C(=O)NR^{15}R^{16}$, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-

~~C₆-C₁₀~~ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 11 of 62

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b},

R^{12b}, at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, aryl, C₃-C₆ cycloalkyl,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

alternatively, R¹⁵ and R¹⁶, together with the nitrogen to which they are attached, may combine to form a 4-7 membered ring wherein said 4-7 membered ring optionally contains an additional heteroatom selected from O or NH;

R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl,
aryl substituted by 0-4 R^{17a}, or
-CH₂-aryl substituted by 0-4 R^{17a},

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 12 of 62

R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃, S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

R¹⁸, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁹, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R^{19b}, at each occurrence, is independently is H or C₁-C₄ alkyl;

R²¹ is H, phenyl, benzyl, or C₁-C₄ alkyl; and

R²² is C₁-C₄ alkyl, C₂-C₄ alkenyl, or C₃-C₄ alkynyl.

2. (CURRENTLY AMENDED) A compound, according to Claim 1, of Formula (I) or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Q is -(CR⁷R^{7a})_m-R⁴,
-(CR⁷R^{7a})_n-S-R⁴,
-(CR⁷R^{7a})_n-O-R⁴, or
-(CR⁷R^{7a})_m-N(R^{7b})-R⁴;

m is 1 or 2;

n is 0 or 1;

R⁴ is H,
C₁-C₈ alkyl substituted with 0-3 R^{4a},
C₂-C₈ alkenyl substituted with 0-3 R^{4a},
C₂-C₈ alkynyl substituted with 0-3 R^{4a},

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 13 of 62

C₃-C₁₀ carbocycle substituted with 0-3 R^{4b},

C₆-C₁₀ aryl substituted with 0-3 R^{4b}, or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b},

R^{4a}, at each occurrence, is independently selected from is H, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, OR^{14a}, C(=O)OR²², SR²², OR²², NR²¹R²², S(=O)R²², S(=O)₂R²²,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,

C₃-C₁₀ carbocycle substituted with 0-3 R^{4b},

C₆-C₁₀ aryl substituted with 0-3 R^{4b}, and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b},

R^{4b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁵ is H;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

C₂-C₆ alkenyl substituted with 0-3 R^{5b};

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)


Page 14 of 62

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};~~C₆-C₁₀~~ aryl substituted with 0-3 R^{5c}; or5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;


R⁶ is H, methyl, or ethyl;R⁷, at each occurrence, is independently H or C₁-C₄ alkyl;R^{7a}, at each occurrence, is independently H or C₁-C₄ alkyl;R^{7b} is H or C₁-C₄ alkyl;

Ring B is selected from:



Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 15 of 62



R^{10} is H , $C(=O)R^{17}$, $C(=O)OR^{17}$, $C(=O)NR^{18}R^{19}$,
 $S(=O)_2NR^{18}R^{19}$, $S(=O)_2R^{17}$;

C_1 - C_6 -alkyl optionally substituted with 0-3 R^{10a} ;

C_6 - C_{10} -aryl substituted with 0-4 R^{10b} ;

C_3 - C_{10} -carbocycle substituted with 0-3 R^{10b} ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{10b} ;

R^{10a} , at each occurrence, is independently selected from H , C_1 - C_6 -alkyl, OR^{14} , Cl , F , Br , I , $=O$, CN , NO_2 , $NR^{15}R^{16}$, CF_3 , or aryl substituted with 0-4 R^{10b} ;

R^{10b} , at each occurrence, is independently selected from H , OH , Cl , F , Br , I , CN , NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, C_1 - C_6 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkoxy, and C_1 - C_4 -haloalkyl S ;

R^{11} , at each occurrence, is independently selected from H , C_1 - C_4 -alkoxy, Cl , F , Br , I , $=O$, CN , NO_2 , $NR^{18}R^{19}$, $C(=O)R^{17}$, $C(=O)OR^{17}$, $C(=O)NR^{18}R^{19}$, $S(=O)_2NR^{18}R^{19}$, CF_3 ;

C_1 - C_6 -alkyl optionally substituted with 0-3 R^{11a} ;

C_6 - C_{10} -aryl substituted with 0-3 R^{11b} ;

C_3 - C_{10} -carbocycle substituted with 0-3 R^{11b} ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b} ;

R^{11a} , at each occurrence, is independently selected from

H , C_1 - C_6 -alkyl, OR^{14} , Cl , F , Br , I , $=O$, CN , NO_2 , $NR^{15}R^{16}$, CF_3 ;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 16 of 62

~~phenyl substituted with 0-3 R^{11b};~~
~~C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and~~
~~5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen,~~
~~oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-~~
~~3 R^{11b};~~

~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂,~~
~~NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,~~
~~C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,~~
~~C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S;~~

W is a bond or -(CH₂)_p;

p is 1 or 2;

X is a bond;

~~phenyl substituted with 0-2 R^{Xb};~~
~~C₃-C₆ carbocycle substituted with 0-2 R^{Xb}; or~~
~~5 to 6 membered heterocycle substituted with 0-2 R^{Xb};~~

~~R^{Xb}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂,~~
~~NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy,~~
~~C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, and C₁-C₃ halothioalkoxy;~~

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -
 NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -
 OC(=O)-;

Z is H;

~~C₁-C₈ alkyl substituted with 0-3 R^{12a},~~
~~C₂-C₆ alkenyl substituted with 0-3 R^{12a},~~
~~C₂-C₆ alkynyl substituted with 0-3 R^{12a},~~
~~C₆-C₁₀ aryl substituted with 0-4 R^{12b},~~
~~C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or~~

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 17 of 62

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b},

R^{12a}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,

C₆-C₁₀ aryl substituted with 0-4 R^{12b},

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b},

R^{12b}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl,

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, benzyl, phenethyl,

(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 18 of 62

alternatively, R¹⁵ and R¹⁶, together with the nitrogen to which they are attached, may combine to form a 4-7 membered ring wherein said 4-7 membered ring optionally contains an additional heteroatom selected from O or NH;

R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, aryl substituted by 0-4 R^{17a}, or -CH₂-aryl substituted by 0-4 R^{17a};

R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃, S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;


R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl;

R^{19b}, at each occurrence, is independently is H or C₁-C₄ alkyl;

R²¹ is H, phenyl, benzyl, or C₁-C₄ alkyl; and

R²² is C₁-C₄ alkyl, C₂-C₄ alkenyl, or C₃-C₄ alkynyl.

3. (CURRENTLY AMENDED) A compound, according to Claim 2, of Formula (Ib):



(Ib)

or a pharmaceutically acceptable salt or prodrug thereof,

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
 BMS PH 7164 (C)
 Page 19 of 62

wherein:

Q is $-(\text{CHR}^7)_m\text{R}^4$,
 $-(\text{CHR}^7)_n\text{S-R}^4$,
 $-(\text{CHR}^7)_n\text{O-R}^4$, or
 $-(\text{CHR}^7)_m\text{N(R}^7b)\text{-R}^4$;

m is 1 or 2;

n is 0 or 1;

R^4 is H,

$\text{C}_1\text{-C}_8$ alkyl substituted with 0-3 R^{4a} ,
 $\text{C}_2\text{-C}_8$ alkenyl substituted with 0-3 R^{4a} ,
 $\text{C}_2\text{-C}_8$ alkynyl substituted with 0-3 R^{4a} ,
 $\text{C}_3\text{-C}_{10}$ carbocycle substituted with 0-3 R^{4b} ,
 ~~$\text{C}_6\text{-C}_{10}$~~ aryl substituted with 0-3 R^{4b} , or
 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b} ;

R^{4a} , at each occurrence, is independently selected from is H, Cl, F, Br, I, CN, NO_2 , $\text{NR}^{15}\text{R}^{16}$, CF_3 , OR^{14a} , C(=O)OR^{22} , SR^{22} , OR^{22} , $\text{NR}^{21}\text{R}^{22}$, S(=O)R^{22} , $\text{S(=O)}_2\text{R}^{22}$, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_1\text{-C}_4$ haloalkyl-S-, $\text{C}_3\text{-C}_{10}$ carbocycle substituted with 0-3 R^{4b} , ~~$\text{C}_6\text{-C}_{10}$~~ aryl substituted with 0-3 R^{4b} , and 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b} ;

R^{4b} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , $\text{NR}^{15}\text{R}^{16}$, CF_3 , acetyl, SCH_3 , S(=O)CH_3 , $\text{S(=O)}_2\text{CH}_3$, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ haloalkyl,

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 20 of 62

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁵ is H;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

C₂-C₆ alkenyl substituted with 0-3 R^{5b};

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

~~C₆-C₁₀~~ aryl substituted with 0-3 R^{5c}; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, Cl, F, Br, I, =O, CN, NO₂, R¹⁵R¹⁶;

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

~~C₆-C₁₀~~ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂,

NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;


R⁷, at each occurrence, is independently H, methyl, or ethyl;

R^{7b} is H, methyl, or ethyl;

Ring B is selected from:

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 21 of 62



~~R¹¹, at each occurrence, is independently selected from~~
~~H, C₁-C₄-alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹, C(=O)R¹⁷, C(=O)OR¹⁷,~~
~~C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃,~~
~~C₁-C₆-alkyl optionally substituted with 0-3 R^{11a},~~
~~C₆-C₁₀-aryl substituted with 0-3 R^{11b},~~
~~C₃-C₁₀-carbocycle substituted with 0-3 R^{11b}, or~~
~~5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen,~~
~~oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with~~
~~0-3 R^{11b},~~

~~R^{11a}, at each occurrence, is independently selected from~~
~~H, C₁-C₆-alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃,~~
~~phenyl substituted with 0-3 R^{11b},~~
~~C₃-C₆-cycloalkyl substituted with 0-3 R^{11b}, and~~
~~5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen,~~
~~oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-~~
~~3 R^{11b},~~

~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂,~~
~~NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,~~
~~C₁-C₆-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl,~~
~~C₁-C₄-haloalkoxy, and C₁-C₄-haloalkyl-S-~~

W is a bond;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 22 of 62

X is a bond;

Y is a bond;

Z is H;

C1-C8 alkyl substituted with 0-3 R^{12a};

C2-C6 alkenyl substituted with 0-3 R^{12a};

C2-C6 alkynyl substituted with 0-3 R^{12a};

C6-C10 aryl substituted with 0-4 R^{12b};

C3-C10 carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C1-C6 alkyl, C1-C4 alkoxy, C1-C4 haloalkyl,

C1-C4 haloalkoxy, C1-C4 haloalkyl-S-,

C6-C10 aryl substituted with 0-4 R^{12b};

C3-C10 carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C1-C6 alkyl, C1-C4 alkoxy, C1-C4 haloalkyl,

C1-C4 haloalkoxy, and C1-C4 haloalkyl-S-;

R¹³, at each occurrence, is independently selected from

H, OH, C1-C6 alkyl, C1-C4 alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C1-C6 alkyl, C2-C6 alkoxyalkyl, or C3-C6 cycloalkyl;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 23 of 62

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl,

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

alternatively, R¹⁵ and R¹⁶, together with the nitrogen to which they are attached, may combine to form a 4-7 membered ring wherein said 4-7 membered ring optionally contains an additional heteroatom selected from O or NH;

R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, aryl substituted by 0-4 R^{17a}, or -CH₂-aryl substituted by 0-4 R^{17a};

R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃, S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl;

R²¹ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl; and

R²² is methyl, ethyl, propyl, butyl, propenyl, butenyl, and propargyl.

4. (CURRENTLY AMENDED) A compound according to Claim 3 of Formula (I)

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 24 of 62

or a pharmaceutically acceptable salt or ~~prodrug~~ thereof,
wherein:

Q is $-(CH_2)_m-R^4$,
 $-(CH_2)_n-S-R^4$,
 $-(CH_2)_n-O-R^4$, or
 $-(CH_2)_m-N(H)-R^4$;

m is 1 or 2;

n is 0 or 1;

R^4 is C_1 - C_8 alkyl substituted with 0-3 R^{4a} ,
 C_2 - C_8 alkenyl substituted with 0-3 R^{4a} ,
 C_2 - C_8 alkynyl substituted with 0-3 R^{4a} ,
 C_3 - C_{10} carbocycle substituted with 0-3 R^{4b} ,
 C_6 - C_{10} aryl substituted with 0-3 R^{4b} , or
 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b} ;

R^{4a} , at each occurrence, is independently selected from is H, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , $C(=O)OR^{22}$, SR^{22} , OR^{22} , OR^{14a} , $NR^{21}R^{22}$, $S(=O)R^{22}$, $S(=O)_2R^{22}$, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 haloalkyl-S-, C_3 - C_{10} carbocycle substituted with 0-3 R^{4b} , C_6 - C_{10} aryl substituted with 0-3 R^{4b} , and 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b} ;

R^{4b} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl,

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
 BMS PH 7164 (C)
 Page 25 of 62

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁵ is H;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

C₂-C₆ alkenyl substituted with 0-3 R^{5b};

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, Cl, F, Br, I, =O, CN, NO₂, R¹⁵R¹⁶;

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};


R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂,

NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and


C₁-C₄ haloalkoxy;

Ring B is selected from:



Serial No. 10/685,031

Response to Office Action of July 8, 2004;
 BMS PH 7164 (C)
 Page 26 of 62



~~R¹¹, at each occurrence, is independently selected from~~

~~H, -O, NR¹⁸R¹⁹, CF₃;~~

~~C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};~~

~~phenyl substituted with 0-3 R^{11b};~~

~~C₃-C₆ carbocycle substituted with 0-3 R^{11b}; and~~

~~5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{11b}; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;~~

~~R^{11a}, at each occurrence, is independently selected from H, C₁-C₄ alkyl, OR¹⁴, F, Cl, -O, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b};~~

~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;~~

W is a bond;

X is a bond;

Y is a bond;

Z is H;

C₁-C₈ alkyl substituted with 0-3 R^{12a};

C₂-C₆ alkenyl substituted with 0-3 R^{12a}; or

C₂-C₆ alkynyl substituted with 0-3 R^{12a};

R^{12a}, at each occurrence, is independently selected from

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 27 of 62

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,
~~C₆-C₁₀~~ aryl substituted with 0-4 R^{12b},
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}, or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b}; and wherein said 5 to 10 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, tetrazolyl, benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl, 1H-indazolyl, oxazolidinyl, isoxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazolinyl, quinolinyl, and isoquinolinyl;

R^{12b}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-; and

alternatively, R¹⁵ and R¹⁶, together with the nitrogen to

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 28 of 62

which they are attached, may combine to form a 4-6 membered ring wherein said 4-6 membered ring optionally contains an additional heteroatom selected from O or NH, wherein said 4-6 membered ring is selected from imidazolidinyl, oxazolidinyl, thiazolidinyl, piperazinyl, morpholinyl, and thiomorpholinyl;

R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl;

R²¹ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl; and

R²² is methyl, ethyl, propyl, butyl, propenyl, butenyl, and propargyl.

5. (CURRENTLY AMENDED) A compound according to Claim 4 wherein:

Q is -CH₂R⁴, -O-R⁴, or -CH₂-NH-R⁴;

R⁴ is C₁-C₆ alkyl substituted with 0-3 R^{4a},
C₂-C₆ alkenyl substituted with 0-3 R^{4a},
C₂-C₆ alkynyl substituted with 0-3 R^{4a},
C₃-C₆ carbocycle substituted with 0-3 R^{4b},
phenyl substituted with 0-3 R^{4b}, or
5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{4b};

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 29 of 62

R^{4a}, at each occurrence, is independently selected from H, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C(=O)OR²², SR²², OR^{14a}, OR²², NR²¹R²², S(=O)R²², S(=O)₂R²², C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-, C₃-C₁₀ carbocycle substituted with 0-3 R^{4b}, C₆-C₁₀ aryl substituted with 0-3 R^{4b}, and 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b};

R^{4b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;


R⁵ is H;
C₁-C₆ alkyl substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b}; or
C₂-C₆ alkynyl substituted with 0-3 R^{5b};

R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, CF₃, Cl, F, Br, I, =O;
C₃-C₆ carbocycle substituted with 0-3 R^{5c};
phenyl substituted with 0-3 R^{5c}; or
5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

Ring B is selected from:

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 30 of 62~~R¹¹, at each occurrence, is independently selected from~~~~H, =O, NR¹⁸R¹⁹, CF₃;~~~~C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};~~~~phenyl substituted with 0-3 R^{11b};~~~~C₃-C₆ carbocycle substituted with 0-3 R^{11b}; and~~~~5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{11b}; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;~~~~R^{11a}, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, phenoxy, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b};~~~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;~~

W is a bond;

X is a bond;

Y is a bond;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 31 of 62

Z is H;

C₁-C₄ alkyl substituted with 0-3 R^{12a};

C₂-C₄ alkenyl substituted with 0-3 R^{12a}; or

C₂-C₄ alkynyl substituted with 0-3 R^{12a};

R^{12a}, at each occurrence, is independently selected from

H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

R¹³, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₄ alkyl, and benzyl;

R¹⁶, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)₂-, and ethyl-S(=O)₂-;

R¹⁸, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl;

R¹⁹, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, and butyl;

R²¹ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl; and

R²² is methyl, ethyl, propyl, butyl, propenyl, butenyl, and propargyl.

6. (CURRENTLY AMENDED) A compound according to Claim 5 or a pharmaceutically acceptable salt or ~~pro-drug~~ thereof wherein:

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 32 of 62

Q is $-\text{CH}_2\text{R}^4$, $-\text{O}-\text{R}^4$, or $-\text{CH}_2-\text{NH}-\text{R}^4$;

R^4 is C_1 - C_6 alkyl substituted with 0-2 R^{4a} ,
 C_2 - C_6 alkenyl substituted with 0-2 R^{4a} ,
 C_2 - C_6 alkynyl substituted with 0-2 R^{4a} , or
 C_3 - C_6 cycloalkyl substituted with 0-3 R^{4b} ;

R^{4a} , at each occurrence, is independently selected from is H, OH, F, Cl, Br, I, CN, $\text{NR}^{15}\text{R}^{16}$, CF_3 , methyl, ethyl, propyl, methoxy, ethoxy, propoxy, OCF_3 ;
 C_3 - C_6 carbocycle substituted with 0-3 R^{4b} ,
 phenyl substituted with 0-3 R^{4b} , or
 5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{4b} ; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R^{4b} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , $\text{NR}^{15}\text{R}^{16}$, CF_3 , acetyl, SCH_3 , $\text{S}(=\text{O})\text{CH}_3$, $\text{S}(=\text{O})_2\text{CH}_3$, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-;

R^5 is H;

C_1 - C_4 alkyl substituted with 0-1 R^{5b} ;
 C_2 - C_4 alkenyl substituted with 0-1 R^{5b} ; or
 C_2 - C_4 alkynyl substituted with 0-1 R^{5b} ;

R^{5b} , at each occurrence, is independently selected from:

 H, methyl, ethyl, propyl, butyl, CF_3 ;
 C_3 - C_6 carbocycle substituted with 0-2 R^{5c} ;
 phenyl substituted with 0-3 R^{5c} ; and
 5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-

Serial No. 10/685,031

Response to Office Action of July 8, 2004;


BMS PH 7164 (C)

Page 33 of 62

3 R^{5c}; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

Ring B is selected from:



~~R¹¹, at each occurrence, is independently selected from~~

~~H, -O, NR¹⁸R¹⁹,~~

~~C₁-C₄ alkyl optionally substituted with 0-1 R^{11a},~~

~~phenyl substituted with 0-3 R^{11b},~~

~~5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{11b}, wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;~~

~~R^{11a}, at each occurrence, is independently selected from H, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, phenoxy, F, Cl, -O, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b},~~

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 34 of 62

~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;~~

W is a bond;

X is a bond;

Y is a bond;

Z is H;

C₁-C₄ alkyl substituted with 0-1 R^{12a};

C₂-C₄ alkenyl substituted with 0-1 R^{12a}; or

C₂-C₄ alkynyl substituted with 0-1 R^{12a};

R^{12a}, at each occurrence, is independently selected from

H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

R¹³, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, Cl, F, Br, CN, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R¹⁵, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl; and

R¹⁶, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, benzyl, and phenethyl;

R¹⁸, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

R¹⁹, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, and butyl.

7. (CURRENTLY AMENDED) A compound according to Claim 6 wherein:

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
 BMS PH 7164 (C)
 Page 35 of 62

R⁵ is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₃,
 -CH₂CH₂CH₂CH₂CH₃, -CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH₂CH₃,
 -CH₂CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH(CH₃)₂, -CH₂NH₂,
 -CH₂N(CH₃)₂, -CH₂N(CH₂CH₃)₂, -CH₂CH₂NH₂, -CH₂CH₂N(CH₃)₂,
 -CH₂CH₂N(CH₂CH₃)₂, -CH₂-cyclopropyl, -CH₂-cyclobutyl,
 -CH₂-cyclopentyl, -CH₂-cyclohexyl,
 -CH₂CH₂-cyclopropyl, -CH₂CH₂-cyclobutyl,
 -CH₂CH₂-cyclopentyl, or -CH₂CH₂-cyclohexyl;

Q is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -
 CH₂CH₂CH₂CH₂CH₃, -CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH₂CH₃, -
 CH₂CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH₂CH₂CH₃,
 -CH₂CH₂CH₂CH₂CH(CH₃)₂, -CH₂-cyclopropyl,
 -CH₂-cyclobutyl, -CH₂-cyclopentyl, -CH₂-cyclohexyl,
 -CH₂CH₂-cyclopropyl, -CH₂CH₂-cyclobutyl,
 -CH₂CH₂-cyclopentyl, -CH₂CH₂-cyclohexyl,
 -OCH₃, -OCH₂CH₃, -OCH₂CH₂CH₃, -OCH(CH₃)₂,
 -OCH₂CH₂CH₂CH₃, -OCH₂CH(CH₃)₂, -OCH₂CH₂CH(CH₃)₂,
 -OCH₂CH₂CH₂CH₂CH₃, -OCH₂CH₂CH₂CH₂CH₂CH₃,
 -OCH₂CH₂CH₂CH(CH₃)₂, -OCH₂CH₂CH₂CH₂CH(CH₃)₂,
 -OCH₂-cyclopropyl, -OCH₂-cyclobutyl,
 -OCH₂-cyclopentyl, -OCH₂-cyclohexyl,
 -OCH₂CH₂-cyclopropyl, -OCH₂CH₂-cyclobutyl,
 -OCH₂CH₂-cyclopentyl, -OCH₂CH₂-cyclohexyl,
 -CH₂OCH₂CH₃, -CH₂OCH₂CH₂CH₃, -CH₂-OCH(CH₃)₂,
 -CH₂OCH₂CH₂CH₂CH₃, -CH₂OCH₂CH(CH₃)₂,
 -CH₂OCH₂CH₂CH₂CH₂CH₃, -CH₂OCH₂CH₂CH(CH₃)₂,
 -CH₂OCH₂CH₂CH₂CH(CH₃)₂, -CH₂O-cyclopropyl,
 -CH₂O-cyclobutyl, -CH₂O-cyclopentyl,
 -CH₂O-cyclohexyl, -CH₂OCH₂-cyclopropyl,
 -CH₂OCH₂-cyclobutyl, -CH₂OCH₂-cyclopentyl,
 -CH₂OCH₂-cyclohexyl, -CH₂(NH)CH₃,
 -CH₂(NH)CH₂CH₃, -CH₂(NH)CH₂CH₂CH₃, -CH₂-(NH)CH(CH₃)₂,
 -CH₂(NH)CH₂CH₂CH₂CH₃, -CH₂(NH)CH₂CH(CH₃)₂,

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 36 of 62

-CH₂(NH)CH₂CH₂CH₂CH₂CH₃, -CH₂(NH)CH₂CH₂CH(CH₃)₂,
 -CH₂(NH)CH₂CH₂CH₂CH(CH₃)₂, -CH₂(NH)-cyclopropyl,
 -CH₂(NH)-cyclobutyl, -CH₂(NH)-cyclopentyl,
 -CH₂(NH)-cyclohexyl, -CH₂(NH)CH₂-cyclopropyl,
 -CH₂(NH)CH₂-cyclobutyl, -CH₂(NH)CH₂-cyclopentyl,
 or -CH₂(NH)CH₂-cyclohexyl;

W is a bond;

X is a bond;

Y is a bond;

Z is methyl, ethyl, i-propyl, n-propyl, n-butyl, i-butyl, s-butyl, t-butyl, or allyl;

~~R¹¹, at each occurrence, is independently selected from~~

~~H, -O, methyl, ethyl, phenyl, benzyl, phenethyl,
 4-F phenyl, (4-F phenyl)CH₂, (4-F phenyl)CH₂CH₂,
 3-F phenyl, (3-F phenyl)CH₂, (3-F phenyl)CH₂CH₂,
 2-F phenyl, (2-F phenyl)CH₂, (2-F phenyl)CH₂CH₂,
 4-Cl phenyl, (4-Cl phenyl)CH₂, (4-Cl phenyl)CH₂CH₂,
 3-Cl phenyl, (3-Cl phenyl)CH₂, (3-Cl phenyl)CH₂CH₂,
 4-CH₃-phenyl, (4-CH₃-phenyl)CH₂, (4-CH₃-phenyl)CH₂CH₂,
 3-CH₃-phenyl, (3-CH₃-phenyl)CH₂, (3-CH₃-phenyl)CH₂CH₂,
 4-CF₃-phenyl, (4-CF₃-phenyl)CH₂, (4-CF₃-phenyl)CH₂CH₂,
 pyrid-2-yl, 4-F pyrid-2-yl, 4-Cl pyrid-2-yl,
 4-CH₃-pyrid-2-yl, 4-CF₃-pyrid-2-yl, pyrid-3-yl,
 4-F pyrid-3-yl, 4-Cl pyrid-3-yl, 4-CH₃-pyrid-3-yl,
 4-CF₃-pyrid-3-yl, or pyrid-4-yl; and~~

~~R¹³, at each occurrence, is independently selected from~~

~~H, F, Cl, OH, -CH₃, -CH₂CH₃, -OCH₃, or -CF₃.~~

8. (CURRENTLY AMENDED) A compound according to Claim 2 of Formula (I) or a pharmaceutically acceptable salt or prodrug thereof

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 37 of 62

wherein:

Q is $-(CH_2)_m-R^4$,
 $-(CH_2)_n-S-R^4$,
 $-(CH_2)_n-O-R^4$, or
 $-(CH_2)_m-N(H)-R^4$;

m is 1 or 2;

n is 0 or 1;

R^4 is C₁-C₈ alkyl substituted with 0-3 R^{4a} ,
 C₂-C₈ alkenyl substituted with 0-3 R^{4a} ,
 C₂-C₈ alkynyl substituted with 0-3 R^{4a} ,
 C₃-C₁₀ carbocycle substituted with 0-3 R^{4b} ,
~~C₆-C₁₀~~ aryl substituted with 0-3 R^{4b} , or
 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b} ;

R^{4a} , at each occurrence, is independently selected from is H, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C(=O)OR²², SR²², OR²², OR^{14a}, NR²¹R²², S(=O)R²², S(=O)₂R²², C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-, C₃-C₁₀ carbocycle substituted with 0-3 R^{4b} , ~~C₆-C₁₀~~-aryl substituted with 0-3 R^{4b} , and 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{4b} ;

R^{4b} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

Serial No. 10/685,031


Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 38 of 62


 R^5 is H;C₁-C₆ alkyl substituted with 0-3 R^{5b} ;C₂-C₆ alkenyl substituted with 0-3 R^{5b} ;C₂-C₆ alkynyl substituted with 0-3 R^{5b} ;C₃-C₁₀ carbocycle substituted with 0-3 R^{5c} ;C₆-C₁₀ aryl substituted with 0-3 R^{5c} ; and5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c} ; R^{5b} , at each occurrence, is independently selected from:H, C₁-C₆ alkyl, CF₃, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶;C₃-C₁₀ carbocycle substituted with 0-3 R^{5c} ;C₆-C₁₀ aryl substituted with 0-3 R^{5c} ; or~~5 to 10~~ membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c} ; R^{5c} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂,NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, andC₁-C₄ haloalkoxy;

Ring B is selected from:



Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 39 of 62



~~R¹¹, at each occurrence, is independently selected from~~

~~H, -O, NR¹⁸R¹⁹, CF₃;~~

~~C₁-C₄ alkyl optionally substituted with 0-3 R^{11a};~~

~~phenyl substituted with 0-3 R^{11b};~~

~~C₃-C₆ carbocycle substituted with 0-3 R^{11b}; or~~

~~5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{11b}, and wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;~~

~~R^{11a}, at each occurrence, is independently selected from~~

~~H, C₁-C₄ alkyl, OR¹⁴, Cl, F, -O, CN, NO₂, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b};~~

~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₄ haloalkoxy;~~

W is a bond, -CH₂-, -CH₂CH₂-;

X is a bond;

phenyl substituted with 0-2 R^{Xb};

C₃-C₆ cycloalkyl substituted with 0-2 R^{Xb}; or

5 to 6 membered heterocycle substituted with 0-2 R^{Xb};

Serial No. 10/685,031

Response to Office Action of July 8, 2004;

BMS PH 7164 (C)

Page 40 of 62

R^{Xb} , at each occurrence, is independently selected from H, OH, Cl, F, $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

Y is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-N(R^{19})-$, $-C(=O)NR^{19b}-$, $-NR^{19b}C(=O)-$, $-NR^{19b}S(=O)_2-$, $-S(=O)_2NR^{19b}-$, $-NR^{19b}S(=O)-$, $-S(=O)NR^{19b}-$, $-C(=O)O-$, or $-OC(=O)-$;

Z is C₁-C₃ alkyl substituted with 1-2 R^{12a} ,
C₆-C₁₀ aryl substituted with 0-4 R^{12b} ,
C₃-C₁₀ carbocycle substituted with 0-3 R^{12b} ; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b} ;

R^{12a} , at each occurrence, is independently selected from
C₆-C₁₀ aryl substituted with 0-4 R^{12b} ,
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b} ; and
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b} ;

R^{12b} , at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and
C₁-C₄ haloalkyl-S-;

R^{13} , at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, and CF_3 ;

R^{14} is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

*Serial No. 10/685,031**Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 41 of 62*

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-; and

alternatively, R¹⁵ and R¹⁶, together with the nitrogen to which they are attached, may combine to form a 4-6 membered ring wherein said 4-6 membered ring optionally contains an additional heteroatom selected from O or NH, wherein said 4-6 membered ring is selected from imidazolidinyl, oxazolidinyl, thiazolidinyl, piperazinyl, morpholinyl, and thiomorpholinyl;

R¹⁸, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁹, at each occurrence, is independently selected from
H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl;

R²¹ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl; and

R²² is methyl, ethyl, propyl, butyl, propenyl, butenyl, and propargyl.

9. (CURRENTLY AMENDED) A compound according to Claim 8 wherein:

Q is -CH₂R⁴, -O-R⁴, or -CH₂-NH-R⁴;

R⁴ is C₁-C₆ alkyl substituted with 0-3 R^{4a},
C₂-C₆ alkenyl substituted with 0-3 R^{4a};

*Serial No. 10/685,031**Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 42 of 62*

C₂-C₆ alkynyl substituted with 0-3 R^{4a},
C₃-C₆ carbocycle substituted with 0-3 R^{4b},
phenyl substituted with 0-3 R^{4b}, or
5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen,
oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-
3 R^{4b},

R^{4a}, at each occurrence, is independently selected from H, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶,
CF₃, C(=O)OR²², SR²², OR^{14a}, OR²², NR²¹R²², S(=O)R²², S(=O)₂R²²,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,
C₃-C₁₀ carbocycle substituted with 0-3 R^{4b},
~~C₆-C₁₀~~-aryl substituted with 0-3 R^{4b}, and
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen,
oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with
0-3 R^{4b},

R^{4b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂,
NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-,

R⁵ is H;

C₁-C₆ alkyl substituted with 0-3 R^{5b},
C₂-C₆ alkenyl substituted with 0-3 R^{5b}, or
C₂-C₆ alkynyl substituted with 0-3 R^{5b},


R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, CF₃, Cl, F, Br, I, =O;
C₃-C₆ carbocycle substituted with 0-3 R^{5c},
phenyl substituted with 0-3 R^{5c}, or
5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen,
oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-
3 R^{5c},

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 43 of 62

R^{5c} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

Ring B is selected from:



~~R^{11} , at each occurrence, is independently selected from H, -O, NR¹⁸R¹⁹, CF₃, C₁-C₄ alkyl optionally substituted with 0-3 R^{11a} , phenyl substituted with 0-3 R^{11b} , C₃-C₆ carbocycle substituted with 0-3 R^{11b} , or 5- to 6-membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5- to 6-membered heterocycle is substituted with 0-3 R^{11b} , and wherein said 5- to 6-membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;~~

~~R^{11a} , at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, phenoxy, Cl, F, -O, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b} ;~~

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 44 of 62

~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₄ haloalkoxy;~~

W is a bond, -CH₂-, -CH₂CH₂-;

X is a bond;

phenyl substituted with 0-1 R^{Xb};

C₃-C₆ cycloalkyl substituted with 0-1 R^{Xb}; or

5 to 6 membered heterocycle substituted with 0-1 R^{Xb};

R^{Xb} is selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, and -OCF₃;

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-,
-N(CH₃)-, or -N(CH₂CH₃)-;

Z is C₁-C₂ alkyl substituted with 1-2 R^{12a};

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from

*Serial No. 10/685,031**Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 45 of 62*

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₄ alkyl, and benzyl;

R¹⁶, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)₂-, and ethyl-S(=O)₂-;

R¹⁸, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl;

R¹⁹, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, and butyl; and

R²¹ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl; and

R²² is methyl, ethyl, propyl, butyl, propenyl, butenyl, and propargyl.

10. (CURRENTLY AMENDED) A compound according to Claim 9 or a pharmaceutically acceptable salt or pro-drug thereof wherein:

Q is -CH₂R⁴, -O-R⁴, or -CH₂-NH-R⁴;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 46 of 62

R⁴ is C₁-C₆ alkyl substituted with 0-2 R^{4a},
C₂-C₆ alkenyl substituted with 0-2 R^{4a},
C₂-C₆ alkynyl substituted with 0-2 R^{4a}, or
C₃-C₆ cycloalkyl substituted with 0-3 R^{4b},

R^{4a}, at each occurrence, is independently selected from is H, OH, F, Cl, Br, I, CN, NR¹⁵NR¹⁶, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, OCF₃;
C₃-C₆ carbocycle substituted with 0-3 R^{4b},
phenyl substituted with 0-3 R^{4b}, or
5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{4b}; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R^{4b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-

R⁵ is H;
C₁-C₄ alkyl substituted with 0-1 R^{5b},
C₂-C₄ alkenyl substituted with 0-1 R^{5b}, or
C₂-C₄ alkynyl substituted with 0-1 R^{5b},

R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, CF₃;
C₃-C₆ carbocycle substituted with 0-2 R^{5c},
phenyl substituted with 0-3 R^{5c}; and
5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{5c}; wherein said 5 to 6 membered heterocycle is selected from pyridinyl,


Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 47 of 62

pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

Ring B is selected from:



~~R¹¹, at each occurrence, is independently selected from~~

~~H, -O, NR¹⁸R¹⁹,~~

~~C₁-C₄ alkyl optionally substituted with 0-3 R^{11a},~~

~~phenyl substituted with 0-3 R^{11b},~~

~~C₃-C₆ carbocycle substituted with 0-3 R^{11b}, or~~

~~5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{11b}, and wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;~~

~~R^{11a}, at each occurrence, is independently selected from~~

~~H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, phenoxy, Cl, F, -O, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b},~~

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 48 of 62

~~R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;~~

W is a bond or -CH₂-;

X is a bond;

phenyl substituted with 0-1 R^{Xb};

C₃-C₆ cycloalkyl substituted with 0-1 R^{Xb}; or

5 to 6 membered heterocycle substituted with 0-1 R^{Xb};

R^{Xb} is selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, methyl, ethyl, methoxy, ethoxy, and -OCF₃;

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-,
-N(CH₃)-, or -N(CH₂CH₃)-;

Z is C₁-C₂ alkyl substituted with 1-2 R^{12a};

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b}; and wherein said 5 to 10 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, tetrazolyl, benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl, 1H-indazolyl, oxazolidinyl, isoxazolidinyl,

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 49 of 62

benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazoliny, quinolinyl, and isoquinolinyl;

R^{12b}, at each occurrence, is independently selected from

H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, and -OCF₃;

R¹³, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, Cl, F, Br, CN, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R¹⁵, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl; and

R¹⁶, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, benzyl, and phenethyl;

R¹⁸, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

R¹⁹, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, and butyl.

11. (CURRENTLY AMENDED) A compound, according to Claim 10, wherein:

R⁵ is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₃,
-CH₂CH₂CH₂CH₂CH₃, -CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH₂CH₃,
-CH₂CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH(CH₃)₂, -CH₂NH₂,
-CH₂N(CH₃)₂, -CH₂N(CH₂CH₃)₂, -CH₂CH₂NH₂, -CH₂CH₂N(CH₃)₂,
-CH₂CH₂N(CH₂CH₃)₂, -CH₂-cyclopropyl, -CH₂-cyclobutyl,
-CH₂-cyclopentyl, -CH₂-cyclohexyl,
-CH₂CH₂-cyclopropyl, -CH₂CH₂-cyclobutyl,
-CH₂CH₂-cyclopentyl, or -CH₂CH₂-cyclohexyl;

Serial No. 10/685,031


Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 50 of 62

Q is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₃, -
CH₂CH₂CH₂CH₂CH₃, -CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH₂CH₃, -
CH₂CH₂CH₂CH(CH₃)₂, -CH₂CH₂CH₂CH₂CH₂CH₂CH₃, -
-CH₂CH₂CH₂CH₂CH(CH₃)₂, -CH₂-cyclopropyl,
-CH₂-cyclobutyl, -CH₂-cyclopentyl, -CH₂-cyclohexyl,
-CH₂CH₂-cyclopropyl, -CH₂CH₂-cyclobutyl,
-CH₂CH₂-cyclopentyl, -CH₂CH₂-cyclohexyl,
-OCH₃, -OCH₂CH₃, -OCH₂CH₂CH₃, -OCH(CH₃)₂,
-OCH₂CH₂CH₂CH₃, -OCH₂CH(CH₃)₂, -OCH₂CH₂CH(CH₃)₂,
-OCH₂CH₂CH₂CH₂CH₃, -OCH₂CH₂CH₂CH₂CH₂CH₃,
-OCH₂CH₂CH₂CH(CH₃)₂, -OCH₂CH₂CH₂CH₂CH(CH₃)₂,
-OCH₂-cyclopropyl, -OCH₂-cyclobutyl,
-OCH₂-cyclopentyl, -OCH₂-cyclohexyl,
-OCH₂CH₂-cyclopropyl, -OCH₂CH₂-cyclobutyl,
-OCH₂CH₂-cyclopentyl, -OCH₂CH₂-cyclohexyl,
-CH₂OCH₂CH₃, -CH₂OCH₂CH₂CH₃, -CH₂-OCH(CH₃)₂,
-CH₂OCH₂CH₂CH₂CH₃, -CH₂OCH₂CH(CH₃)₂,
-CH₂OCH₂CH₂CH₂CH₂CH₃, -CH₂OCH₂CH₂CH(CH₃)₂,
-CH₂OCH₂CH₂CH₂CH(CH₃)₂, -CH₂O-cyclopropyl,
-CH₂O-cyclobutyl, -CH₂O-cyclopentyl,
-CH₂O-cyclohexyl, -CH₂OCH₂-cyclopropyl,
-CH₂OCH₂-cyclobutyl, -CH₂OCH₂-cyclopentyl,
-CH₂OCH₂-cyclohexyl; -CH₂(NH)CH₃,
-CH₂(NH)CH₂CH₃, -CH₂(NH)CH₂CH₂CH₃, -CH₂-(NH)CH(CH₃)₂,
-CH₂(NH)CH₂CH₂CH₂CH₃, -CH₂(NH)CH₂CH(CH₃)₂,
-CH₂(NH)CH₂CH₂CH₂CH₂CH₃, -CH₂(NH)CH₂CH₂CH(CH₃)₂,
-CH₂(NH)CH₂CH₂CH₂CH(CH₃)₂, -CH₂(NH)-cyclopropyl,
-CH₂(NH)-cyclobutyl, -CH₂(NH)-cyclopentyl,
-CH₂(NH)-cyclohexyl, -CH₂(NH)CH₂-cyclopropyl,
-CH₂(NH)CH₂-cyclobutyl, -CH₂(NH)CH₂-cyclopentyl,
or -CH₂(NH)CH₂-cyclohexyl;

W is a bond or -CH₂-;

X is a bond;

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 51 of 62

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, or -N(CH₃)-,

Z is phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl, 2-Cl-phenyl, 3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl, 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl, 3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl, 2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl, 3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl, 3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl, 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl, 4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl, 2-CF₃O-phenyl, 3-CF₃O-phenyl, 4-CF₃O-phenyl, furanyl, thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl, 4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl, 1-benzimidazolyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, morpholino, N-piperinyl, phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-, (4-F-phenyl)CH₂-, (2-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂-, (2,3-diF-phenyl)CH₂-, (2,4-diF-phenyl)CH₂-, (2,5-diF-phenyl)CH₂-, (2,6-diF-phenyl)CH₂-, (3,4-diF-phenyl)CH₂-, (3,5-diF-phenyl)CH₂-, (2,3-diCl-phenyl)CH₂-, (2,4-diCl-phenyl)CH₂-, (2,5-diCl-phenyl)CH₂-, (2,6-diCl-phenyl)CH₂-, (3,4-diCl-phenyl)CH₂-, (3,5-diCl-phenyl)CH₂-, (3-F-4-Cl-phenyl)CH₂-, (3-F-5-Cl-phenyl)CH₂-, (3-Cl-4-F-phenyl)CH₂-, (2-MeO-phenyl)CH₂-, (3-MeO-phenyl)CH₂-, (4-MeO-phenyl)CH₂-, (2-Me-phenyl)CH₂-,

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 52 of 62

(3-Me-phenyl)CH₂-, (4-Me-phenyl)CH₂-,
(2-MeS-phenyl)CH₂-, (3-MeS-phenyl)CH₂-,
4-MeS-phenyl)CH₂-, (2-CF₃O-phenyl)CH₂-,
(3-CF₃O-phenyl)CH₂-, (4-CF₃O-phenyl)CH₂-,
(furanlyl)CH₂-, (thienyl)CH₂-, (pyridyl)CH₂-,
(2-Me-pyridyl)CH₂-, (3-Me-pyridyl)CH₂-,
(4-Me-pyridyl)CH₂-, (1-imidazolyl)CH₂-,
(oxazolyl)CH₂-, (isoxazolyl)CH₂-,
(1-benzimidazolyl)CH₂-, (cyclopropyl)CH₂-, (cyclobutyl)CH₂-, (cyclopentyl)CH₂-,
(cyclohexyl)CH₂-, (morpholino)CH₂-,
(N-piperidinyl)CH₂-, or (phenyl)₂CH-;

~~R¹¹~~, at each occurrence, is independently selected from

~~H, -O, methyl, ethyl, phenyl, benzyl, phenethyl,
4-F-phenyl, (4-F-phenyl)CH₂-, (4-F-phenyl)CH₂CH₂-,
3-F-phenyl, (3-F-phenyl)CH₂-, (3-F-phenyl)CH₂CH₂-,
2-F-phenyl, (2-F-phenyl)CH₂-, (2-F-phenyl)CH₂CH₂-,
4-Cl-phenyl, (4-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂CH₂-,
3-Cl-phenyl, (3-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂CH₂-,
4-CH₃-phenyl, (4-CH₃-phenyl)CH₂-, (4-CH₃-phenyl)CH₂CH₂-,
3-CH₃-phenyl, (3-CH₃-phenyl)CH₂-, (3-CH₃-phenyl)CH₂CH₂-,
4-CF₃-phenyl, (4-CF₃-phenyl)CH₂-, (4-CF₃-phenyl)CH₂CH₂-,
pyrid-2-yl, 4-F-pyrid-2-yl, 4-Cl-pyrid-2-yl,
4-CH₃-pyrid-2-yl, 4-CF₃-pyrid-2-yl, pyrid-3-yl,
4-F-pyrid-3-yl, 4-Cl-pyrid-3-yl, 4-CH₃-pyrid-3-yl,
4-CF₃-pyrid-3-yl, or pyrid-4-yl; and~~


R¹³, at each occurrence, is independently selected from

H, F, Cl, OH, -CH₃, -CH₂CH₃, -OCH₃, or -CF₃.

12. (CURRENTLY AMENDED) A compound according to Claim 2 ~~one of Claims 4-11~~ of Formula (Ic):

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 53 of 62



(Ic)

or a stereoisomer, pharmaceutically acceptable salt ~~or prodrug~~ thereof.

13.-16. (CANCELLED)

17. (CURRENTLY AMENDED) A compound according to Claim 1, or a pharmaceutically acceptable salt ~~or prodrug~~ thereof comprising:

(7S)-[(2S)-1-oxo-2-pentyloxy-4-methylpentyl]amino-5-methyl-5H,7H-dibenzo[b,d]azepin-6-one.

18. (ORIGINAL) A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

19.-20 (CANCELLED)

21. (NEW) A pharmaceutical composition comprising a compound of Claim 2 and a pharmaceutically acceptable carrier.

22. (NEW) A pharmaceutical composition comprising a compound of Claim 3 and a pharmaceutically acceptable carrier.

23. (NEW) A pharmaceutical composition comprising a compound of Claim 4 and a pharmaceutically acceptable carrier.

24. (NEW) A pharmaceutical composition comprising a compound of Claim 5 and a pharmaceutically acceptable carrier.

25. (NEW) A pharmaceutical composition comprising a compound of Claim 6 and a pharmaceutically acceptable carrier.

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 54 of 62

26. (NEW) A pharmaceutical composition comprising a compound of Claim 7 and a pharmaceutically acceptable carrier.
27. (NEW) A pharmaceutical composition comprising a compound of Claim 8 and a pharmaceutically acceptable carrier.
28. (NEW) A pharmaceutical composition comprising a compound of Claim 9 and a pharmaceutically acceptable carrier.
29. (NEW) A pharmaceutical composition comprising a compound of Claim 10 and a pharmaceutically acceptable carrier.
30. (NEW) A pharmaceutical composition comprising a compound of Claim 11 and a pharmaceutically acceptable carrier.
31. (NEW) A pharmaceutical composition comprising a compound of Claim 12 and a pharmaceutically acceptable carrier.
32. (NEW) A pharmaceutical composition comprising a compound of Claim 17 and a pharmaceutically acceptable carrier.
33. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 1.
34. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 2.
35. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 3.
36. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 4.

Serial No. 10/685,031

Response to Office Action of July 8, 2004;
BMS PH 7164 (C)
Page 55 of 62

37. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 5.
38. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 6.
39. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 7.
40. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 8.
41. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 9.
42. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 10.
43. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 11.
44. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 12.
45. (NEW) A method for the treatment of Alzheimer's Disease comprising administering to a host in need of such treatment a therapeutically effective amount of a compound of Claim 17.